arXiv:hep-ph/9403343v2 6 May 1994

HUPD-9406
hep-ph/9403343
March, 1994

Effective Lagrangian for a Technicolor Model without Exact Custodial Symmetry

T.Yoshikawa, H.Takata and T.Morozumi

Department of Physics, Hiroshima University
1-3-1 Kagamiyama, Higashi-Hiroshima, 724
Japan

Abstract

Effective Lagrangian including technimesons is constructed for a realistic one-family Technicolor model without exact custodial symmetry. Tree level contribution to oblique correction parameters S and U due to spin 1 technimesons are computed with the effective Lagrangian. An isospin breaking term which is associated with technilepton vector mesons gives a negative contribution to the electroweak radiative correction parameter S due to mixing between $I = 0$ and $I = 1$ vector mesons. U receives non-zero contribution due to exotic left-handed charged vector mesons and its sign can be both negative and positive.

1 Introduction

It has been shown that Technicolor models are strongly constrained by precision measurements of electorweak parameters. In particular, QCD scale up one-family Technicolor model with exact custodial symmetry seem to be already excluded by studying an oblique correction parameter S [1].

$$S_{theory} = 0.28 \times 4 = 1.1$$

$$S_{exp} = -0.42 \pm 0.45$$

where S_{theory} is an estimation with vector and axial vector meson dominance assumption for QCD scale up one-family Technicolor model. (See appendix E.) A factor of 4 comes from the fact that the model contains 4 $SU(2)$ doublet of technifermions. S_{exp} is quoted from [2] and the reference point of the standard model is taken at $m_t = 150GeV$ and $m_H = 1TeV$. However, according to ref.[3], this is not the case for Technicolor models without exact custodial symmetry. A realistic model is proposed for a one-family Technicolor model. In their model, isospin breaking is introduced for a light technilepton doublet. The doublet contributes to a radiative correction parameter S in negative sign while keeping ρ parameter nearly equal to 1. In their analysis, free technifermion model is used to compute S parameter.

In this paper, we compute the oblique corrections (S, T and U) in an alternative way. We construct an effective Lagrangian with low lying technimesons for the one-family Technicolor model without exact custodial symmetry. By using the effective Lagrangian, we can compute non-perturbative effect on the radiative correction parameters S,T, and U due to bound states of technifermions. Our paper is organized in the following way. In §2, we review feature of the model. With some assumption on the low lying technimesons' spectrum, we construct a low energy effective Lagrangian. In §3, S, T, and U parameters are computed with the Lagrangian. It is shown that S parameter receives negative contribution due to the mixing between isosinglet and isotriplet techni-vector mesons. §4 is devoted to finding the range of the paramerters for negative S. Conclusions and discussion are summarized in §5.

2 Effective Lagrangian for a Technicolor model without exact custodial symmetry

Let us describe the model briefly [3]. The model has a global symmetry : $G = SU(6)_L \otimes SU(6)_R \otimes SU(2)_L \otimes U(1)_{2R} \otimes U(1)_{8L} \otimes U(1)_{8R} \otimes U(1)_V$ which is spontaneously broken to $H = SU(6)_V \otimes U(1)_{2V} \otimes U(1)_{8V} \otimes U(1)_V$. The part of G must be broken explicitly in order that unnecessary massless physical Nambu-Goldstone bosons disappear.

The technifermions are assigned to the following representations of $SU(3)_c \otimes SU(2) \otimes U(1)_Y \otimes SU(N_{TC})$:

$$\begin{aligned}
(U,D)_L &= (3,2,\frac{Y_{Lq}}{2},N_{TC}),\\
U_R &= (3,1,\frac{Y_{Lq}}{2}+\frac{1}{2},N_{TC}),\\
D_R &= (3,1,\frac{Y_{Lq}}{2}-\frac{1}{2},N_{TC}),\\
(N,E)_L &= (1,2,\frac{Y_{Ll}}{2}+\frac{1}{2},N_{TC}),\\
E_R &= (1,1,\frac{Y_{Ll}}{2}-\frac{1}{2},N_{TC}),
\end{aligned}$$

where $Y_{Lq}(Y_{Ll})$ is hypercharge of lefthanded techniquark(technilepton) ($Y_{Lq} = 1/3$, $Y_{Ll} = -1$).

The following mass spectrum is assumed for technifermions:

(1) $M_U = M_D$,

(2) $M_N < M_E < M_U$.

$SU(6)_V$ symmetry is preserved because techniquarks are degenerate, while $SU(2)_V$ symmetry is explicitly broken due to isospin breaking of technileptons. To proceed further, we need to know the technimesons spectrum of the model. Since the model does not have the same global symmetry as that of QCD ($SU(2)_L \otimes SU(2)_R$ in chiral limit) and underlying dynamics may be also different from that, we can not simply scale up the mesons' spectrum in QCD. Here we make use of the global symmetry as a guide to construct an effective Lagrangian. Global symmetry strongly constraints the structure of effective Lagrangian as well as properties of bound states included in effective Lagrangian. Concerned with technimesons, which are bound states of technifermions, we need to make a few assumptions. In this paper, we include only spin 0 (Nambu Goldstone Bosons (NGB) and Pseudo Nambu Goldstone Bosons (PNGB)) and spin 1 mesons. For the purpose of studying tree level contribution to the oblique correction parameters, S,T, and U, other mesons with higher spins (spin≥ 2) can be ignored because they do not contribute to self energy corrections of gauge bosons. Further we only keep $O(p^2)$ terms for NGB and PNGB sector and ignore their loop effects and $O(p^4)$ counter terms. About spin 1 mesons, we employ the approach of including vector mesons into chiral Lagrangian [4][5] and extend it to our case. In table(1), technimesons and their technifermion contents as well as their J,P,C, and I are listed.(For spin 0 sector, we quote them from ref.[3].) Note that charged technilepton NGBs ($\Pi^{\pm}$) do not have definite parity because $\Pi^{\pm}$ are NGBs associated with $SU(2)_L$ not $SU(2)_A$. (Note that we do not have full $SU(2)_R$ symmetry.) In the same way, exotic left-handed charged vector mesons(A_L) are introduced so that they interact with $\Pi^{\pm}$ and $SU(2) \otimes U(1)$ gauge bosons etc without loss of the invariance. Corresponding to spin0, vector, and axial and left-handed vector mesons, the effective Lagrangian consists of three parts: $\mathcal{L}_S$, $\mathcal{L}_V$, and $\mathcal{L}_A$: The explicit form for them will be presented below.

$\mathcal{L}_S$ (spin0 sector)

This part of the Lagrangian consists of $O(p^2)$ terms of NGBs and PNGBs.

$$\mathcal{L}_S = F_6^2 tr(\alpha_{6\perp})^2 + F_L^2 tr(\alpha_{L\perp})^2 + F_2^2 tr(\alpha_{2\perp})^2 + F_8^2 tr(\alpha_{8\perp})^2$$

$$+ \quad \beta_{\perp} tr(\alpha_{8\perp}\tau^3\alpha_{2\perp}) + \mathcal{L}'_{br}, \tag{1}$$

where,

$$\begin{aligned}
\alpha_{6\perp\mu} &= 2\sum_{\alpha=1}^{35}\begin{pmatrix} T^{\alpha} & \\ & 0 \end{pmatrix} tr[T^{\alpha}\frac{\xi_6\nabla_{R\mu}\xi_6^{\dagger} - \xi_6^{\dagger}\nabla_{L\mu}\xi_6}{2i}], \qquad (2)\\
\alpha_{L\perp\mu} &= -2\sum_{a=1}^{2}\frac{1}{2}\begin{pmatrix} 0 & \\ & \tau^{a} \end{pmatrix} tr[\frac{\tau^a}{2}\frac{\xi_2^{\dagger}\hat{\nabla}_{L\mu}\xi_2}{2i}], \qquad (3)\\
\alpha_{2\perp\mu} &= 2\frac{1}{2}\begin{pmatrix} 0 & \\ & \tau^{3} \end{pmatrix} tr[\frac{\tau^3}{2}\frac{\xi_1^{\dagger}\hat{\nabla}_{R\mu}\xi_1 - \xi_2^{\dagger}\hat{\nabla}_{L\mu}\xi_2}{2i}], \qquad (4)\\
\alpha_{8\perp\mu} &= 2\frac{1}{4\sqrt{3}}\begin{pmatrix} I_6 & \\ & -3I_2 \end{pmatrix} [tr\{\frac{1}{4\sqrt{3}}\frac{\xi_6\nabla_{R\mu}\xi_6^{\dagger} - \xi_6^{\dagger}\nabla_{L\mu}\xi_6}{2i}\}\\
&\qquad + tr\{\frac{-3}{4\sqrt{3}}\frac{\xi_1^{\dagger}\hat{\nabla}_{R\mu}\xi_1 - \xi_2^{\dagger}\hat{\nabla}_{L\mu}\xi_2}{2i}\}], \qquad (5)
\end{aligned}$$

where,

$$\begin{aligned}
\nabla_{R\mu} &= \partial_\mu + ig'(\frac{\tau^3}{2}\otimes I_3)B_\mu + ig'(\frac{Y_{LQ}}{2}\otimes I_3)B_\mu, \qquad (6)\\
\nabla_{L\mu} &= \partial_\mu + ig\sum_{A=1}^{3}(\frac{\tau^A}{2}\otimes I_3)W_\mu^A + ig'(\frac{Y_{LQ}}{2}\otimes I_3)B_\mu, \qquad (7)\\
\hat{\nabla}_{R\mu} &= \partial_\mu + ig'\frac{\tau^3}{2}B_\mu + ig'\frac{Y_{Ll}}{2}B_\mu, \qquad (8)\\
\hat{\nabla}_{L\mu} &= \partial_\mu + ig\sum_{A=1}^{3}\frac{\tau^A}{2}W_\mu^A + ig'\frac{Y_{Ll}}{2}B_\mu, \qquad (9)
\end{aligned}$$

$$\begin{aligned}
\xi_6 &= \exp(i\sum_{\alpha=1}^{35}\frac{T^\alpha P^\alpha}{F_6} + i\frac{1}{4\sqrt{3}}\frac{\theta_8}{F_8}), \qquad (10)\\
\xi_2 &= \exp(i\sum_{a=1}^{2}\frac{\tau^a\Pi^a}{2F_L} + i\frac{\tau^3\Pi^3}{2F_2} - i\frac{3}{4\sqrt{3}}\frac{\theta_8}{F_8}), \qquad (11)\\
\xi_1 &= \exp(-i\frac{\tau^3\Pi^3}{2F_2} + i\frac{3}{4\sqrt{3}}\frac{\theta_8}{F_8}). \qquad (12)
\end{aligned}$$

In (1) $\mathcal{L}'_{br}$ consists of explicit breaking terms which make physical NGB massive. Without $\mathcal{L}'_{br}$ and $SU(2)_L\otimes U(1)_Y$ gauge interaction, we have 3 color singlet physical massless NGBs which are linear combinations of $\theta_8, \Pi^3, \Pi^{\pm}, P^3$ and $P^{\pm}$. Here $P^{\pm}$ and P^3 are $SU(2)$ triplet and color singlet bound states of techniquarks. By adding $\mathcal{L}'_{br}$, we can make them massive. (See appendix D for the details.) In (3), τ^a is a projection into $SU(2)_L$ $(a = 1, 2)$ part. In (6)-(9), W^A and B are $SU(2) \otimes U(1)_Y$ gauge bosons. In (10)-(12), ξs correspond to NGB fields. T^α are generator of $SU(6)$ and τ^a, τ^3 are Pauli matrices. P^α are 35 NGBs for broken $SU(6)_A$ symmetry, which are techniquark bound states. $\Pi^a (a = 1, 2)$ and Π^3 are NGBs for broken $SU(2)_L(\tau^a : a = 1, 2)$ and $U(1)_{2A}$ symmetry respectively, which are technilepton bound states. θ_8 is a NGB for $U(1)_{8A}$ symmetry. F_6, F_L, F_2 and F_8 are decay constants for these NGBs. Note that F_L is not degenerate with F_2 because custodial symmetry is broken in the sector and Π_L form an irreducible representation under $U(1)_{2V}$ which Π^3 does not belong to. The difference between F_L and F_2 gives rise to small deviation of ρ parameter from 1 (see (76)). The transformation of ξs are given by;

$$\xi'_6 = g_{L6}\xi_6 h_6^\dagger = h_6 \xi_6 g_{R6}^\dagger, \tag{13}$$

$$\xi'_2 = g_{L2}\xi_2 h_2^\dagger, \tag{14}$$

$$\xi'_1 = g_{R2}\xi_1 h_2^\dagger, \tag{15}$$

$$\frac{\theta'_8}{F_8} = \frac{\theta_8}{F_8} + \phi_{R8} - \phi_{L8}, \tag{16}$$

where,

$$g_{L6} = \exp\,(iT_6^\alpha \phi_{L6}^\alpha + i\frac{1}{4\sqrt{3}}\phi_{L8} + i\frac{1}{4}\phi_V), \tag{17}$$

$$g_{R6} = \exp\,(iT_6^\alpha \phi_{R6}^\alpha + i\frac{1}{4\sqrt{3}}\phi_{R8} + i\frac{1}{4}\phi_V), \tag{18}$$

$$g_{L2} = \exp\,(i\sum_{a=1}^{2} \tau^a \phi_{L2}^a + i\tau^3 \phi_{L2}^3 - i\frac{3}{4\sqrt{3}}\phi_{L8} + i\frac{1}{4}\phi_V), \tag{19}$$

$$g_{R2} = \exp\,(i\tau^3 \phi_{R2}^3 - i\frac{3}{4\sqrt{3}}\phi_{R8} + i\frac{1}{4}\phi_V), \tag{20}$$

$$h_6 = \hat{h}_6 h_{Q1}, \tag{21}$$

$$\hat{h}_6 = \exp(iT_6^\alpha \phi_{V6}^\alpha), \tag{22}$$

$$h_{Q1} = \exp(i\frac{1}{2\sqrt{3}}\phi_{Q1}), \tag{23}$$

$$h_2 = \hat{h}_2 h_{L1}, \tag{24}$$

$$\hat{h}_2 = \exp(i\frac{\tau^3}{2}\phi_{V2}), \tag{25}$$

$$h_{L1} = \exp(i\frac{1}{2}\phi_{L1}), \tag{26}$$

and ϕ_{L8} and ϕ_{R8} are x independent parameters of $U(1)_{8L} \otimes U(1)_{8R}$ transformations. It is not difficult to see that $\alpha_{6\perp\mu}$, $\alpha_{L\perp\mu}$ and $\alpha_{2\perp\mu}$ transform as:

$$\alpha'_{6\perp\mu} = \hat{h}_6 \alpha_{6\perp\mu} \hat{h}_6^\dagger, \tag{27}$$

$$\alpha'_{2L\mu} = \hat{h}_2 \alpha_{2L\mu} \hat{h}_2^\dagger, \tag{28}$$

$$\alpha'_{2\perp\mu} = \alpha_{2\perp\mu}, \tag{29}$$

$$\alpha'_{8\perp\mu} = \alpha_{8\perp\mu}. \tag{30}$$

$\mathcal{L}$ (Vector meson(1^{--}) sector)

In addition to NGBs and PNGBs, we incorporate vector mesons into the effective Lagrangian. Corresponding to unbroken symmetry: $SU(6)_V \otimes U(1)_{2V} \otimes U(1)_{8V} \otimes U(1)_V$, we introduce 38 vector mesons, 35 of them are techniquark bound states which belong to the adjoint representation of $SU(6)_V$. Corresponding to $U(1)_{qV}$ ($U(1)_V$ for techniquark sector) ,$U(1)_{lV}$ ($U(1)_V$ for technilepton sector) and $U(1)_{2V}$, three neutral vector mesons, *techni* ω_q ($\omega_{6\mu}$), *techni* ω_l ($\omega_{2\mu}$), and *techni* ρ_l ($\rho_{2\mu}$) are introduced. For the techniquark sector of the effective lagrangian, we can just extend the approach of [4] [5] into the larger symmetry, i.e, chiral $SU(6)_L \otimes SU(6)_R \otimes U(1)_{6V}$. On the otherhand, for technilepton sector, non-trivial isospin breaking terms are introduced. Let us record vector meson part of the effective Lagrangian first,

$$\begin{aligned}
\mathcal{L}_V &= \frac{1}{2}trF^{\mu\nu}_{V_6}F_{V_6\,\mu\nu}+\frac{1}{2}trF^{\mu\nu}_{V_2}F_{V_2\,\mu\nu}+\frac{1}{2}trF^{\mu\nu}_{V_{\omega 6}}F_{V_{\omega 6}\,\mu\nu}+\frac{1}{2}trF^{\mu\nu}_{V_{\omega 2}}F_{V_{\omega 2}\,\mu\nu} \\
&- M^2_{V_6}tr(V_{6\mu}-\frac{i}{G_6}\alpha_{6\|\mu})^2 - M^2_{V_{\omega 6}}tr(V_{\omega 6\mu}-\frac{i}{G_{\omega 6}}\alpha_{\omega 6\|\mu})^2 \\
&- M^2_{V_2}tr(V_{2\mu}-\frac{i}{G_2}\alpha_{2\|\mu})^2 - M^2_{V_{\omega 2}}tr(V_{\omega 2\mu}-\frac{i}{G_{\omega 2}}\alpha_{\omega 2\|\mu})^2 \\
&+ \alpha_V trF^{\mu\nu}_{V_2}\tau^3F_{V_{\omega 2}\,\mu\nu} \\
&- \beta_V tr(V_{2\mu}-\frac{i}{G_2}\alpha_{2\|\mu})\tau^3(V_{\omega 2\mu}-\frac{i}{G_{\omega 2}}\alpha_{\omega 2\|\mu}),
\end{aligned} \tag{31}$$

where

$$\alpha_{6\|_\mu} = 2\sum_{\alpha=1}^{35}\begin{pmatrix}T^\alpha & \\ & 0\end{pmatrix}tr[T^\alpha\frac{\xi_6\nabla_{R\mu}\xi_6^\dagger+\xi_6^\dagger\nabla_{L\mu}\xi_6}{2i}], \tag{32}$$

$$\alpha_{\omega 6\|_\mu} = 2\frac{1}{2\sqrt{3}}\begin{pmatrix}I_6 & \\ & 0\end{pmatrix}tr[\frac{1}{2\sqrt{3}}\frac{\xi_6\nabla_{R\mu}\xi_6^\dagger+\xi_6^\dagger\nabla_{L\mu}\xi_6}{2i}], \tag{33}$$

$$\alpha_{2\|_\mu} = 2\frac{1}{2}\begin{pmatrix}0 & \\ & \tau^3\end{pmatrix}tr[\frac{\tau^3}{2}\frac{\xi_1^\dagger\hat{\nabla}_{R\mu}\xi_1+\xi_2^\dagger\hat{\nabla}_{L\mu}\xi_2}{2i}], \tag{34}$$

$$\alpha_{\omega 2\|_\mu} = 2\frac{1}{2}\begin{pmatrix}0 & \\ & I_2\end{pmatrix}tr[\frac{1}{2}\frac{\xi_1^\dagger\hat{\nabla}_{R\mu}\xi_1+\xi_2^\dagger\hat{\nabla}_{L\mu}\xi_2}{2i}]. \tag{35}$$

Vector mesons are decomposed into their component fields,

$$V_{6\mu} = i\sum_{\alpha=1}^{35}\begin{pmatrix}T^\alpha & \\ & 0\end{pmatrix}\rho^\alpha_{6\mu}, \tag{36}$$

$$V_{\omega 6\mu} = i\frac{1}{2\sqrt{3}}\begin{pmatrix}I_6 & \\ & 0\end{pmatrix}\omega_{6\mu}, \tag{37}$$

$$V_{2\mu} = i\frac{1}{2}\begin{pmatrix}0 & \\ & \tau^3\end{pmatrix}\rho_{2\mu}, \tag{38}$$

$$V_{\omega 2\mu} = i\frac{1}{2}\begin{pmatrix}0 & \\ & I_2\end{pmatrix}\omega_{2\mu}. \tag{39}$$

The quantities defined above transform under G in the following way,

$$\begin{cases}\alpha'_{6\|\mu} &= \hat{h}_6\alpha_{6\|\mu}\hat{h}_6^\dagger - i\hat{h}_6\partial_\mu\hat{h}_6^\dagger \\ V'_{6\mu} &= \hat{h}_6V_{6\mu}\hat{h}_6^\dagger+\frac{1}{G_6}\hat{h}_6\partial_\mu\hat{h}_6^\dagger.\end{cases}, \tag{40}$$

$$\begin{cases} \alpha'_{\omega6\|\mu} & = \alpha_{\omega6\|\mu} - ih_{Q1}\partial_\mu h^\dagger_{Q1} \\ V'_{\omega6\mu} & = V_{\omega6\mu} + \frac{1}{G_{\omega6}} h_{Q1}\partial_\mu h^\dagger_{Q1}. \end{cases}, \tag{41}$$

$$\begin{cases} \alpha'_{2\|\mu} & = \alpha_{2\|\mu} - i\hat{h}_2\partial_\mu \hat{h}^\dagger_2 \\ V'_{2\mu} & = V_{2\mu} + \frac{1}{G_{\omega2}} \hat{h}_2\partial_\mu \hat{h}^\dagger_2. \end{cases}, \tag{42}$$

$$\begin{cases} \alpha'_{\omega2\|\mu} & = \alpha_{\omega2\|\mu} - ih_{L1}\partial_\mu h^\dagger_{L1} \\ V'_{\omega2\mu} & = V_{\omega2\mu} + \frac{1}{G_{\omega2}} h_{L1}\partial_\mu h^\dagger_{L1}. \end{cases}. \tag{43}$$

In (31), the terms which are proportional to α_V and β_V break isospin symmetry. The term with the coefficient α_V generates mixing between *techni* ω_l ($\omega_{2\mu}$ $(I = 0)$) and *techni* ρ_l ($\rho_{2\mu}$ $(I = 1)$) through the kinetic term while the term which coefficient is β_V generates the mixing through the mass term.

Axial vector meson (A) and left-handed vector meson (A_L) sector

This part of the Lagrangian are given by,

$$\begin{aligned}
\mathcal{L}_A &= \frac{1}{2} tr F^{\mu\nu}_{A_6} F_{A_6\mu\nu} + \frac{1}{2} tr F^{\mu\nu}_{A_L} F_{A_L\mu\nu} + \frac{1}{2} tr F^{\mu\nu}_{A_2} F_{A_2\mu\nu} + \frac{1}{2} tr F^{\mu\nu}_{A_8} F_{A_8\mu\nu} \\
&- M^2_{A_6} tr(A_{6\mu} - \frac{i}{\lambda_6}\alpha_{6\perp\mu})^2 - M^2_{A_8} tr(A_{8\mu} - \frac{i}{\lambda_8}\alpha_{8\perp\mu})^2 \\
&- M^2_{A_L} tr(A_{L\mu} - \frac{i}{\lambda_L}\alpha_{L\perp\mu})^2 - M^2_{A_2} tr(A_{2\mu} - \frac{i}{\lambda_2}\alpha_{2\perp\mu})^2 \\
&- \alpha_A \frac{2}{\sqrt{3}} tr F^{\mu\nu}_{A_2} \tau^3 F_{A_8\mu\nu} \\
&+ \beta_A \frac{2}{\sqrt{3}} tr(A_{2\mu} - \frac{i(1+\delta)}{\lambda_2}\alpha_{2\perp\mu})\tau^3(A_{8\mu} - \frac{i(1+\delta')}{\lambda_8}\alpha_{8\perp\mu}) \\
&+ \frac{1}{4}\frac{\beta^2_A M^2_{A2}\delta^2}{M^2_{A2}M^2_{A8} - \frac{1}{4}\beta^2_A}\frac{1}{\lambda^2_2} tr(\alpha_{2\perp\mu})^2 \\
&+ \frac{1}{4}\frac{\beta^2_A M^2_{A8}\delta'^2}{M^2_{A2}M^2_{A8} - \frac{1}{4}\beta^2_A}\frac{1}{\lambda^2_8} tr(\alpha_{8\perp\mu})^2 \\
&+ \frac{2}{\sqrt{3}}\frac{\delta\delta'\beta_A M^2_{A2}M^2_{A8}}{M^2_{A2}M^2_{A8} - \frac{1}{4}\beta^2_A}\frac{1}{\lambda_2\lambda_8} tr(\alpha_{2\perp\mu}\tau^3\alpha_{8\perp\mu}).
\end{aligned} \tag{44}$$

The decomposition into component fields is given by,

$$A_{6\mu} = i\sum_{\alpha=1}^{35}\begin{pmatrix} T^\alpha & \\ & 0 \end{pmatrix} a^\alpha_{6\mu}, \tag{45}$$

$$
\begin{aligned}
A_{8\mu} &= i\frac{1}{4\sqrt{3}}\begin{pmatrix} I_6 & \\ & -3I_2 \end{pmatrix} a_{8\mu}, && (46)\\
A_{L\mu} &= i\sum_{a=1}^{2}\frac{1}{2}\begin{pmatrix} 0 & \\ & \tau^a \end{pmatrix} a^a_{L\mu}, && (47)\\
A_{2\mu} &= i\frac{1}{2}\begin{pmatrix} 0 & \\ & \tau^3 \end{pmatrix} a_{2\mu}. && (48)
\end{aligned}
$$

In (46), $\tau^a (a = 1, 2)$ is a projection into $SU(2)_L$. They transform under G in the following way,

$$
\begin{aligned}
A'_{6\mu} &= \hat{h}_6 A_{6\mu} \hat{h}^\dagger_6, && (49)\\
A'_{8\mu} &= A_{8\mu}, && (50)\\
A'_{L\mu} &= \hat{h}_2 A_{L\mu} \hat{h}^\dagger_2, && (51)\\
A'_{2\mu} &= A_{2\mu}. && (52)
\end{aligned}
$$

Note that there are not inhomogeneous terms for the transformation of axial (A) and left-handed (A_L) vector mesons under G. Therefore the mixing terms between these vector mesons and PNGB occur. This effect results in redefinition of the coefficients of $O(p^2)$ terms of NGB and PNGB sector ((1)). In order to avoid the redifinition of the coefficients, we just need to add the appropriate $O(p^2)$ terms in axial and left-handed vector meson sector. The $O(p^2)$ terms of NGBs and PNGBs in (44) are chosen so that the $O(p^2)$ terms in (1) will not be altered after eliminating A and A_L with their equation of motion. (See appendix **D** for the details of the procedure.) Compared with vector meson sector, two additional terms with the coefficients δ and δ' come in (44). Both of them are isospin breaking terms and can contribute to S and U through mixing between axial vector mesons (a_2 and a_8) and PNGBs (Π^3 and θ_8). However, because we focus on the contribution due to vector, axial and left-handed vector mesons only, we ignore the contribution due to these terms and put δ and δ' zero in the following analysis.

3 S,T and U parameter

The electroweak radiative correction parameter S, T and U are defined in terms of the gauge boson self-energy:

$$\begin{aligned} S &= 16\pi \frac{d}{dq^2}[\delta\Pi_{33}(q^2) - \delta\Pi_{3Q}(q^2)] \mid_{q^2=0} \\ &= -16\pi \frac{d}{dq^2}[\delta\Pi_{3Y}(q^2)] \mid_{q^2=0}, & (53) \\ \alpha T &= \frac{g^2+g'^2}{M_Z^2}[\delta\Pi_{11}(0) - \delta\Pi_{33}(0)], & (54) \\ U &= 16\pi \frac{d}{dq^2}[\delta\Pi_{11}(q^2) - \delta\Pi_{33}(q^2)] \mid_{q^2=0} . & (55) \end{aligned}$$

where the $\delta\Pi$ is contribution of beyond standard model.

In order to compute technimesons contribution to S, T and U, we need to expand the effective Lagrangian in terms of their component fields explicitly. Because we only consider their tree level contribution here, it is suffice to keep technicolor singlet and color singlet technimesons in the expansion. We also have done one more simplification. In princple, not only vector, axial and lefthanded-vector mesons, PNGBs can contribute to S and U even in the tree level when isospin is not exact symmetry. For example, PNGB θ_8 can mix with W_3 and B through the isospin breaking term which strength is given by $\beta_\perp$ in (1). Since the mixing term is derivative coupling, the contribution to S is proportional to $\beta_\perp^2/M_{\theta 8}^2$. Here $M_{\theta 8}$ is a mass of $\theta_8$which comes from the explicit breaking of $U(1)_{8A}$. Similarly, Π^3 can contribute to S and U through isospin breaking terms whose coefficients are δ and δ'. Since we have not known the masses of PNGBs and how they mix with each other, we just assumed that they are so heavy or the mixing with gauge bosons are so small that their contribution to S and U are negligible. This amounts to the following simplification in the effective Lagrangian.

$$\beta_\perp = \delta = \delta' = 0. \tag{56}$$

Under this assumption, in tree level, it is spin 1 technimesons that contribute to S and U. In the following, we compute S, T and U in techniquark sector and in technilepton sector respectively. The latter computation will tell us how differently custodial symmetry breaking terms contribute to S compared with techniquark sector where custodial symmetry is exact.

The techniquark sector

In this sector, color singlet and $SU(2)$ triplet part of techni-vector mesons $V_6(1^-)$ and $A_6(1^+)$ contribute to S. Since $SU(2)$ singlet vector mesons such as $V_{\omega 6}$ and A_8 do not couple with W_3, they will not contribute to S. The mixing terms between the vector mesons and gauge bosons are given by :

$$\begin{aligned}\mathcal{L}_{6int} &= -M_{V_6}^2 tr(V_{6\mu} - \frac{i}{G_6}\alpha_{6\|\mu})^2 - M_{A_6}^2 tr(A_{6\mu} - \frac{i}{\lambda_6}\alpha_{6\perp\mu})^2 \\ &= 2M_{V_6}^2[\frac{\rho_6}{2} - \frac{\sqrt{3}}{4G_6}(gW_3 + g'B)]^2 + 2M_{A_6}^2[\frac{a_6}{2} + \frac{\sqrt{3}}{4\lambda_6}(gW_3 - g'B)]^2. \end{aligned} \tag{57}$$

$\rho_6(a_6)$ are color singlet and isotriplet vector(axial vector) mesons. Therefore, these terms induce $W_3 - (\rho_6, a_6)$ and $B - (\rho_6, a_6)$ mixing and contributes to S through the Feynman diagram (Fig.(1)). We obtain S in techniquark sector.

$$\begin{aligned} igg'S_q = &-16\pi\frac{d}{dq^2}[(i\sqrt{3}\frac{M_{V6}^2}{2G_6}g)\frac{-i}{q^2 - M_{V6}^2}(ig'\sqrt{3}\frac{M_{V6}^2}{2G_6})]\mid_{q^2=0} \\ &-16\pi\frac{d}{dq^2}[(i\sqrt{3}\frac{M_{A6}^2}{2\lambda_6}g)\frac{-i}{q^2 - M_{A6}^2}(-ig'\sqrt{3}\frac{M_{A6}^2}{2\lambda_6})]\mid_{q^2=0} . \end{aligned} \tag{58}$$

$$S_q = 4\pi[\frac{3}{G_6^2} - \frac{3}{\lambda_6^2}]. \tag{59}$$

The technilepton sector

In this sector, there are isospin breaking terms,

$$\begin{aligned}
\mathcal{L}' &= \alpha_V tr F^{\mu\nu}_{V_2} \tau^3 F_{V_\omega 2_{\mu\nu}} \\
&- \beta_V tr (V_2 - \frac{i}{G_2} \alpha_{2\|\mu}) \tau^3 (V_{\omega 2} - \frac{i}{G_{\omega 2}} \alpha_{\omega 2\|\mu}) \\
&- \alpha_A \frac{2}{\sqrt{3}} tr F^{\mu\nu}_{A_2} \tau^3 F_{A_8\,\mu\nu} \\
&+ \beta_A \frac{2}{\sqrt{3}} tr (A_{2\mu} - \frac{i}{\lambda_2} \alpha_{2\perp\mu}) \tau^3 (A_{8\mu} - \frac{i}{\lambda_8} \alpha_{8\perp\mu}). \qquad (60)
\end{aligned}$$

Because of these terms, isospin of vector mesons is not a conserved quantity and mixing between I=0 and I=1 vector mesons can occur. Then, $SU(2)$ singlet vector mesons such as *techni* ω_l (ω_2) can contribute to S through the mixing terms. Therefore , we need to diagonalize mass terms and kinetic terms of vector mesons. By expanding the effective Lagrangian, we obtain:

$$\begin{aligned}
\mathcal{L}_V &= -\frac{1}{4} \begin{pmatrix} \partial_{[\mu,}\rho_{2\nu]} & \partial_{[\mu,}\omega_{2\nu]} \end{pmatrix} \begin{pmatrix} 1 & \alpha_V \\ \alpha_V & 1 \end{pmatrix} \begin{pmatrix} \partial_{[\mu,}\rho_{2\nu]} \\ \partial_{[\mu,}\omega_{2\nu]} \end{pmatrix} \\
&+\frac{1}{2} \begin{pmatrix} \rho_2 & \omega_2 \end{pmatrix} \begin{pmatrix} M^2_{V2} & \frac{\beta_V}{2} \\ \frac{\beta_V}{2} & M_{V_{\omega 2}} \end{pmatrix} \begin{pmatrix} \rho_2 \\ \omega_2 \end{pmatrix} \\
&-\frac{1}{2} \begin{pmatrix} \rho_2 & \omega_2 \end{pmatrix} \begin{pmatrix} M^2_{V2} & \frac{\beta_V}{2} \\ \frac{\beta_V}{2} & M^2_{V_{\omega 2}} \end{pmatrix} \begin{pmatrix} \frac{gW_3+g'B}{G_2} \\ \frac{2Y_{Ll}g'B}{G_{\omega 2}} \end{pmatrix}, \qquad (61)
\end{aligned}$$

$$\begin{aligned}
\mathcal{L}_A &= -\frac{1}{4} \begin{pmatrix} \partial_{[\mu,}a_{2\nu]} & \partial_{[\mu,}a_{8\nu]} \end{pmatrix} \begin{pmatrix} 1 & \alpha_A \\ \alpha_A & 1 \end{pmatrix} \begin{pmatrix} \partial_{[\mu,}a_{2\nu]} \\ \partial_{[\mu,}a_{8\nu]} \end{pmatrix} \\
&+\frac{1}{2} \begin{pmatrix} a_2 & a_8 \end{pmatrix} \begin{pmatrix} M^2_{A2} & \frac{\beta_A}{2} \\ \frac{\beta_A}{2} & M_{A_8} \end{pmatrix} \begin{pmatrix} a_2 \\ a_8 \end{pmatrix} \\
&+\frac{1}{2} \begin{pmatrix} a_2 & a_8 \end{pmatrix} \begin{pmatrix} M^2_{A2} & \frac{\beta_A}{2} \\ \frac{\beta_A}{2} & M^2_{A_8} \end{pmatrix} \begin{pmatrix} \frac{gW_3-g'B}{\lambda_2} \\ 0 \end{pmatrix}, \qquad (62)
\end{aligned}$$

where

$$\partial_{[\mu,}V_{\nu]} = \partial_\mu V_\nu - \partial_\nu V_\mu.$$

The kinetic and mass terms can be diagonalized by doing the following transformations successively.

$$\begin{pmatrix} \rho_2^m \\ \omega_2^m \end{pmatrix} = U_{mV} Z_V U_{DV} \begin{pmatrix} \rho_2 \\ \omega_2 \end{pmatrix}, \tag{63}$$

$$\begin{pmatrix} a_2^m \\ a_8^m \end{pmatrix} = U_{mA} Z_A U_{DA} \begin{pmatrix} a_2 \\ a_8 \end{pmatrix}, \tag{64}$$

where U_D,Z and U_m are defined by,

$$\begin{aligned} U_{DV(A)} &= \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}, & (65) \\ Z_{V(A)} &= \begin{pmatrix} (1-\alpha_{V(A)})^{\frac{1}{2}} & 0 \\ 0 & (1+\alpha_{V(A)})^{\frac{1}{2}} \end{pmatrix}, & (66) \\ U_{mV(A)} &= \begin{pmatrix} c_{V(A)} & -s_{V(A)} \\ s_{V(A)} & c_{V(A)} \end{pmatrix}. & (67) \end{aligned}$$

U_D is a 45 degree rotation matrix to diagonalize the kinetic terms. Z is a scale transformation to keep correct normalization for the kinetic terms. Finally U_m is a rotation matrix which diagonalizes the mass terms. U_m relates the mass matrices to their eigen values in the following way.

$$\begin{aligned} \begin{pmatrix} M_\rho^2 & 0 \\ 0 & M_\omega^2 \end{pmatrix} &= U_{mV} Z_V^{-1} U_{DV} \begin{pmatrix} M_{V2}^2 & \frac{\beta_V}{2} \\ \frac{\beta_V}{2} & M_{V\omega 2} \end{pmatrix} U_{DV}^{-1} Z_V^{-1} U_{mV}^{-1} \\ = U_{mV} & \begin{pmatrix} \frac{1}{2(1-\alpha_V)}(M_{V2}^2 + M_{V\omega 2}^2 - \beta_V) & \frac{1}{2\sqrt{(1-\alpha_V^2)}}(M_{V2}^2 - M_{V\omega 2}^2) \\ \frac{1}{2\sqrt{(1-\alpha_V^2)}}(M_{V2}^2 - M_{V\omega 2}^2) & \frac{1}{2(1+\alpha_V)}(M_{V2}^2 + M_{V\omega 2}^2 + \beta_V) \end{pmatrix} U_{mV}^{-1}, \end{aligned} \tag{68}$$

$$\begin{aligned} \begin{pmatrix} M_{a2}^2 & 0 \\ 0 & M_{a8}^2 \end{pmatrix} &= U_{mA} Z_A^{-1} U_{DA} \begin{pmatrix} M_{A2}^2 & \frac{\beta_A}{2} \\ \frac{\beta_A}{2} & M_{A8} \end{pmatrix} U_{DA}^{-1} Z_A^{-1} U_{mA}^{-1} \\ = U_{mA} & \begin{pmatrix} \frac{1}{2(1-\alpha_A)}(M_{A2}^2 + M_{A8}^2 - \beta_A) & \frac{1}{2\sqrt{(1-\alpha_A^2)}}(M_{A2}^2 - M_{A8}^2) \\ \frac{1}{2\sqrt{(1-\alpha_A^2)}}(M_{A2}^2 - M_{A8}^2) & \frac{1}{2(1+\alpha_A)}(M_{A2}^2 + M_{A8}^2 + \beta_A) \end{pmatrix} U_{mA}^{-1}. \end{aligned} \tag{69}$$

In (68) and (69), M_ρ and M_ω (M_{a2} and M_{a8}) are eigenvalues of vector (axial vector) mesons mass matrix. With these transformations, the interaction term between gauge bosons and vector mesons are given by:

$$\begin{aligned}\mathcal{L}_{int} \quad = \quad & -\frac{1}{2}\begin{pmatrix} \rho_2^m & \omega_2^m \end{pmatrix}\begin{pmatrix} M_\rho^2 & 0 \\ 0 & M_\omega^2 \end{pmatrix} U_{mV} Z_V U_{DV} \begin{pmatrix} \frac{gW_3+g'B}{G_2} \\ \frac{2Y_{Ll}g'B}{G_{\omega 2}} \end{pmatrix} \\ & +\frac{1}{2}\begin{pmatrix} a_2^m & a_8^m \end{pmatrix}\begin{pmatrix} M_{a2}^2 & 0 \\ 0 & M_{a8}^2 \end{pmatrix} U_{mA} Z_A U_{DA} \begin{pmatrix} \frac{gW_3-g'B}{\lambda_2} \\ 0 \end{pmatrix},\end{aligned} \tag{70}$$

where,

$$\begin{aligned}&\begin{pmatrix} M_\rho^2 & 0 \\ 0 & M_\omega^2 \end{pmatrix} U_{mV} Z_V U_{DV} \\ &= \frac{1}{\sqrt{2}}\begin{pmatrix} M_\rho^2[c_V(1-\alpha_V)^{\frac{1}{2}} - s_V(1+\alpha_V)^{\frac{1}{2}}] & M_\rho^2[-c_V(1-\alpha_V)^{\frac{1}{2}} - s_V(1+\alpha_V)^{\frac{1}{2}}] \\ M_\omega^2[c_V(1+\alpha_V)^{\frac{1}{2}} + s_V(1-\alpha_V)^{\frac{1}{2}}] & M_\omega^2[c_V(1+\alpha_V)^{\frac{1}{2}} - s_V(1-\alpha_V)^{\frac{1}{2}}] \end{pmatrix},\end{aligned} \tag{71}$$

$$\begin{aligned}&\begin{pmatrix} M_{a2}^2 & 0 \\ 0 & M_{a8}^2 \end{pmatrix} U_{mA} Z_A U_{DA} \\ &= \frac{1}{\sqrt{2}}\begin{pmatrix} M_{a2}^2[c_A(1-\alpha_A)^{\frac{1}{2}} - s_A(1+\alpha_A)^{\frac{1}{2}}] & M_{a2}^2[-c_A(1-\alpha_A)^{\frac{1}{2}} - s_A(1+\alpha_A)^{\frac{1}{2}}] \\ M_{a8}^2[c_A(1+\alpha_A)^{\frac{1}{2}} + s_A(1-\alpha_A)^{\frac{1}{2}}] & M_{a8}^2[c_A(1+\alpha_A)^{\frac{1}{2}} - s_A(1-\alpha_A)^{\frac{1}{2}}] \end{pmatrix}.\end{aligned} \tag{72}$$

By computing the Feynman diagram (Fig.2), the sum of the contribution to S due to technilepton sector and techniquark sector ((59)) is,

$$S \quad = \quad 4\pi[\frac{3}{G_6^2} - \frac{3}{\lambda_6^2} + \frac{1}{G_2^2} - \frac{1}{\lambda_2^2} + Y_{Ll}\frac{2\alpha_V}{G_2 G_{\omega 2}}]. \tag{73}$$

The term which is proportional to Y_{Ll} is due to isospin breaking. The minimum of S can be obtained for $\alpha_V = 1$ because of $Y_{Ll} = -1$. Only ω_2^m which consists of $N\bar{N}$ component contributes to S in that case, because $E\bar{E}$ component of vector meson decouples and ideal mixing is realized. This confirms that the conjecture of the importance of ω which consists of $N\bar{N}$ in the same model[3]. For T parameter, we obtain the same expression as that is given in ref.[3]. For completeness, we give the explicit form here.

$$M_W^2 = \frac{1}{4}g^2\{3F_6^2 + F_L^2\}, \tag{74}$$

$$M_Z^2 = \frac{1}{4}(g^2 + {g'}^2)\{3F_6^2 + F_2^2\}, \tag{75}$$

$$\alpha T = \rho - 1 = \frac{F_L^2 - F_2^2}{3F_6^2 + F_2^2}. \tag{76}$$

With the assumption:$F_L, F_2 \ll F_6$, T parameter can be so small even if there is a splitting between F_L and F_2 as stated by the authors of [3]. Finally we compute U parameter. U parameter is zero if the isospin symmetry is exact. Therefore the contribution to U in the present model comes only from technilepton sector. The left-handed vector meson a_L contributes to $\delta\Pi_{11}$ part of U (Fig.(3)). This contribution is not cancelled by $\delta\Pi_{33}$ part which $I = 1$ vector (ρ_2) and axial vector (a_2) mesons contribute to.

$$U = 4\pi[\frac{1}{G_2^2} + \frac{1}{\lambda_2^2} - \frac{1}{\lambda_L^2}]. \tag{77}$$

The sign and the value of U depend on three coupling constants ; G_2, λ_2, and λ_L.

4 The range of parameters for negative S

In this section, we explore the parameter region where S parameter is negative since the present experiment fits favor negative S [2]. If the future experiments constraint on S is improved, we may do more complete analysis. Before studying the results of the model without exact custodial symmetry, let us review the estimation of S_q(the contribution of S in techniquark sector) under the assumption that the techniquark sector is just QCD scale up technicolor theory so that we can get a hint of the order of S which we are arguing. If underlying dynamics of Technicolor for techniquark sector is exactly the same as that of QCD ($N_{TC} = N_C = 3$), we can estimate the parameters of the techniquark sector in the effective Lagrangian by simply scaling up the corresponding parameters. Because S_q depends only on dimensionless coupling constants G_6 and λ_6, we may use the same value for the correspoinding coupling constants which are determined by low energy hadronic processes. We have determined G_6 from $\rho \to \pi\pi$ decay and λ_6 from $a_1(1260) \to \pi\gamma$. This leads to ${G_6}^2 = 31.5$ and ${\lambda_6}^2 = 106$.(See appendix E.)

$$S_q = (0.40 - 0.12) \times 3 = 0.28 \times 3 \tag{78}$$

0.40 comes from the contribution of $I = 1$ vector meson while 0.12 comes from $I = 1$ axial vector meson. A factor of 3 comes from the fact that the techniquark sector contains three SU(2) doublet corresponding to color degree of freedom. This result is consistent with the result with dispersion analysis [1] and the results given by chiral lagrangian with vector resonances. [6]

Now let us turn to the present model. Since we have not known the underlying dynamics of the present model, we do not have any guiding principle to determine the parameters of our model without experimental information. Therefore instead of trying to predict S in our model, we determine the allowed region of the parameters of the effective lagrangian by imposing the present experimentally allowed region for S. Since we have many parameters, we further need to limit ourselves into the small parameter space to draw some definite conclusions. Here we simply assume that S is dominated by only vector mesons and the contribution of axial vector mesons can be ignored. As shown in (78), this is a good approximation in case of the QCD scale up technicolor model. However it is not clear if the assumption still holds even in the present model. Nevertheless, let us proceed further. Under the assumption of vector (1^{--}) dominance, S is given by,

$$\begin{aligned} S &= 4\pi[\frac{3}{G_6^2} + \frac{1}{G_2^2} - \alpha_V \frac{2}{G_2 G_{\omega 2}}] \\ &\geq 4\pi[\frac{3}{G_6^2} + \frac{1}{G_2^2} - \frac{2}{G_2 G_{\omega 2}}]. \end{aligned} \tag{79}$$

Here we have substituted $Y_{Ll} = -1$ in (73) and neglect axial vector contribution. In the second line of (79), the inequality holds because α_V can take its value between -1 and 1 and $\alpha_V = 1$ makes S minimum. Note that the bound for α_V comes from the condition for the positive semi-definitness of the kinetic terms of vector mesons (65). In the follwing we assume that $\alpha_V = 1$ and impose the condition of negative S. The condition for $S \leq 0$ now leads a relation,

$$\frac{G_6}{G_2}(\frac{2G_6}{G_{\omega 2}} - \frac{G_6}{G_2}) \geq 3. \tag{80}$$

This region is shown in Fig.(4) in the parameter space (G_6/G_2 vs $G_6/G_{\omega 2}$). We can get the lower bound for one of the parameter, $G_6/G_{\omega 2}$.

$$\frac{G_6}{G_{\omega 2}} \geq \sqrt{3} \tag{81}$$

Hence $G_{\omega 2}$ must be smaller than G_6 in ordet to make S negative. This means that the coupling strength between *techni* ω_l and gauge bosons are stronger than that between *techni* ρ_q and gauge bosons. Note that the coupling strength between gauge bosons and vector mesons is proportional to $1/G$. Since the negative contribution to S is proportional to Y_{Ll} and Y_{Ll} part of hypercharge interaction couples with *techni* ω_l, strong coupling between *techni* ω_l and hypercharge gauge boson B is preferred to get negative S.

5 Conclusions and Discussion

In this paper, we have constructed an effective Lagrangian for a technicolor model without exact custodial symmetry. By using the Lagrangian, we compute tree level contribution to S and U due to spin 1 technimesons. We have shown that in a realistic one-family model, the *techni* ρ_l and the *techni* ω_l mixing can contribute to S parameter with negative sign. The most important term in our effective Lagrangian is the mixing in the kinetic term, $trF_\rho \tau^3 F_\omega$. S is independent of the coeffcient of the mass mixing term β_V. We also study the condition to make S minimum under the vector meson (1^{--}) dominance. We find that the vector meson consists of $N\bar{N}$ component must be dominant dynamical degree of freedom to make S minimum. ($\alpha_V = 1$) This argument holds as far as hypercharge (Y_{Ll}) for which isospin breaking is introduced is negative. Thus the mechanism for negative S presented in this paper does not work for one doublet model with $Y_L = 0$. This conclusion is consistent with an analysis with a free technifermion model [3]. On the contrary to the present model, we may introduce a small isospin breaking for techniquark sector. In that case, the corresponding parameter of isospin breaking term, α_V must be -1 to make S minimum because hypercharge of techniquark Y_{Lq} is positive. The vector mesons consisits of $D\bar{D}$ component will play major role to make S minimum in that case. We also note that exotic lefthanded charged vector mesons are naturally introduced in our framework. They contribute to U due to the mixing with $W^{\pm}$. U can be both negative and positive depending on the parameters. There are many things to be done in this direction. The relation between our computation and that with a free technifermion model must be clarified. The origin of the isospin breaking must be studied. Also we need to relate the parameters of the effective Lagrangian to more fundamental interaction for example, by modeling technicolor by Nambu Jona-Lasinio model [7].

Acknowledgement

We would like to thank V.A. Miransky for informing us of [3] and thank T. Onogi, K. Hikasa and M. Harada for discussions.

Appendix

In this appendix, we provide some useful formulas needed to derive the results given in the text.

A Effective Lagrangian

$$
\begin{aligned}
\mathcal{L}_S &= F_6^2 tr(\alpha_{6\perp})^2 + F_L^2 tr(\alpha_{L\perp})^2 + F_2^2 tr(\alpha_{2\perp})^2 + F_8^2 tr(\alpha_{8\perp})^2 \\
&+ \beta_\perp tr(\alpha_{8\perp}\tau^3\alpha_{2\perp}),
\end{aligned}
\tag{A.1}
$$

$$
\begin{aligned}
\mathcal{L}_V &= \frac{1}{2} tr F^{\mu\nu}_{V_6} F_{V_6\,\mu\nu} + \frac{1}{2} tr F^{\mu\nu}_{V_2} F_{V_2\,\mu\nu} + \frac{1}{2} tr F^{\mu\nu}_{V_{\omega 6}} F_{V_{\omega 6}\,\mu\nu} + \frac{1}{2} tr F^{\mu\nu}_{V_{\omega 2}} F_{V_{\omega 2}\,\mu\nu} \\
&- M^2_{V_6} tr(V_{6\mu} - \frac{i}{G_6}\alpha_{6\parallel\mu})^2 - M^2_{V_{\omega 6}} tr(V_{\omega 6\mu} - \frac{i}{G_{\omega 6}}\alpha_{\omega 6\parallel\mu})^2 \\
&- M^2_{V_2} tr(V_{2\mu} - \frac{i}{G_2}\alpha_{2\parallel\mu})^2 - M^2_{V_{\omega 2}} tr(V_{\omega 2\mu} - \frac{i}{G_{\omega 2}}\alpha_{\omega 2\parallel\mu})^2 \\
&+ \alpha_V tr F^{\mu\nu}_{V_2} \tau^3 F_{V_{\omega 2}\,\mu\nu} \\
&- \beta_V tr(V_{2\mu} - \frac{i}{G_2}\alpha_{2\parallel\mu})\tau^3(V_{\omega 2\mu} - \frac{i}{G_{\omega 2}}\alpha_{\omega 2\parallel\mu}),
\end{aligned}
\tag{A.2}
$$

$$
\begin{aligned}
\mathcal{L}_A &= \frac{1}{2} tr F^{\mu\nu}_{A_6} F_{A_6\,\mu\nu} + \frac{1}{2} tr F^{\mu\nu}_{A_L} F_{A_L\,\mu\nu} + \frac{1}{2} tr F^{\mu\nu}_{A_2} F_{A_2\,\mu\nu} + \frac{1}{2} tr F^{\mu\nu}_{A_8} F_{A_8\,\mu\nu} \\
&- M^2_{A_6} tr(A_{6\mu} - \frac{i}{\lambda_6}\alpha_{6\perp\mu})^2 - M^2_{A_8} tr(A_{8\mu} - \frac{i}{\lambda_8}\alpha_{8\perp\mu})^2 \\
&- M^2_{A_L} tr(A_{L\mu} - \frac{i}{\lambda_L}\alpha_{L\perp\mu})^2 - M^2_{A_2} tr(A_{2\mu} - \frac{i}{\lambda_2}\alpha_{2\perp\mu})^2 \\
&- \alpha_A \frac{2}{\sqrt{3}} tr F^{\mu\nu}_{A_2} \tau^3 F_{A_8\,\mu\nu} \\
&+ \beta_A \frac{2}{\sqrt{3}} tr(A_{2\mu} - \frac{i(1+\delta)}{\lambda_2}\alpha_{2\perp\mu})\tau^3(A_{8\mu} - \frac{i(1+\delta')}{\lambda_8}\alpha_{8\perp\mu}) \\
&+ \frac{1}{4}\frac{\beta_A^2 M_{A2}^2 \delta^2}{M_{A2}^2 M_{A8}^2 - \frac{1}{4}\beta_A^2}\frac{1}{\lambda_2^2} tr(\alpha_{2\perp\mu})^2 \\
&+ \frac{1}{4}\frac{\beta_A^2 M_{A8}^2 \delta'^2}{M_{A2}^2 M_{A8}^2 - \frac{1}{4}\beta_A^2}\frac{1}{\lambda_8^2} tr(\alpha_{8\perp\mu})^2
\end{aligned}
$$

$$+ \quad \frac{2}{\sqrt{3}} \frac{\delta\delta' \beta_A M_{A2}^2 M_{A8}^2}{M_{A2}^2 M_{A8}^2 - \frac{1}{4}\beta_A^2} \frac{1}{\lambda_2 \lambda_8} tr(\alpha_{2\perp\mu} \tau^3 \alpha_{8\perp\mu}). \qquad (\text{A.3})$$

B Decomposition into fields' components

In terms of fields' components, $\alpha_\perp$s and $\alpha_\parallel$s are given by:

$$\begin{aligned}
\alpha_{6\perp\mu} &= -\frac{1}{2\sqrt{3}F_6}\{\sum_{a=1}^{2} \tau^a (\partial_\mu P^a + \frac{\sqrt{3}F_6}{2} g W_\mu^a) \\
&\qquad + \tau^3 (\partial_\mu P^3 + \frac{\sqrt{3}F_6}{2}(gW_\mu^3 - g'B_\mu))\} \otimes I_3, && (\text{B.1}) \\
\alpha_{L\perp\mu} &= -\frac{1}{2F_L}\sum_{a=1}^{2} \tau^a \{\partial_\mu \hat{\Pi}^a + \frac{F_L}{2} g W_\mu^a\}, \quad \hat{\Pi}^a \equiv \frac{1}{2}\Pi^a, && (\text{B.2}) \\
\alpha_{2\perp\mu} &= -\frac{1}{2F_2}\tau^3 \{\partial_\mu \Pi^3 + \frac{F_2}{2}(gW_\mu^3 - g'B_\mu)\}, && (\text{B.3}) \\
\alpha_{8\perp\mu} &= -\frac{1}{4\sqrt{3}F_8}\begin{pmatrix} I_6 & \\ & -3I_2 \end{pmatrix} \partial_\mu \theta_8, && (\text{B.4}) \\
& && (\text{B.5})
\end{aligned}$$

$$\begin{aligned}
A_{6\mu} - \frac{i}{\lambda_6}\alpha_{6\perp\mu} &= \frac{i}{2\sqrt{3}}\tau^3 \otimes I_3 [a_{6\mu}^3 + \frac{1}{\lambda_6 F_6}\{\partial_\mu P^3 + \frac{\sqrt{3}F_6}{2}(gW_\mu^3 - g'B_\mu)\} \\
&+ \frac{i}{2\sqrt{3}}\sum_{a=1}^{2} \tau^a \otimes I_3 [a_{6\mu}^a + \frac{1}{\lambda_6 F_6}\{\partial_\mu P^a + \frac{\sqrt{3}F_6}{2} g W_\mu^a\}], && (\text{B.6}) \\
A_{L\mu} - \frac{i}{\lambda_L}\alpha_{L\perp\mu} &= \frac{i}{2}\sum_{a=1}^{2} \tau^a [a_{L\mu} + \frac{1}{\lambda_L F_L}\{\partial_\mu \hat{\Pi}^a + \frac{F_L}{2} g W_\mu^a\}], && (\text{B.7}) \\
A_{2\mu} - \frac{i}{\lambda_2}\alpha_{2\perp\mu} &= \frac{i}{2}\tau^3 [a_{2\mu} + \frac{1}{\lambda_2 F_6}\{\partial_\mu \Pi^3 + \frac{F_2}{2}(gW_{3\mu} - g'B_\mu)\}], && (\text{B.8}) \\
A_{8\mu} - \frac{i}{\lambda_8}\alpha_{8\perp\mu} &= \frac{i}{4\sqrt{3}}\begin{pmatrix} I_6 & \\ & -3I_2 \end{pmatrix} [a_{8\mu} + \frac{1}{\lambda_8 F_8}\partial_\mu \theta_8], && (\text{B.9}) \\
V_{6\mu} - \frac{i}{G_6}\alpha_{6\parallel\mu} &= \frac{i}{2\sqrt{3}}\tau^3 \otimes I_3 [\rho_{6\mu}^3 - \frac{\sqrt{3}}{2G_6}(gW_{3\mu} + g'B_\mu)] \\
&\quad - \frac{i}{2\sqrt{3}}\sum_{a=1}^{2} \tau^a \otimes I_3 [\rho_{6\mu}^a - \frac{\sqrt{3}}{2G_6}(gW_\mu^a)], && (\text{B.10})
\end{aligned}$$

$$V_{\omega 6\mu} - \frac{i}{G_{\omega 6}}\alpha_{\omega 6\|\mu} = \frac{i}{2\sqrt{3}} I_6[\omega_{6\mu} - \frac{\sqrt{3}}{2G_{\omega 6}}(2Y_{Ll}g'B_\mu)], \tag{B.11}$$

$$V_{2\mu} - \frac{i}{G_2}\alpha_{2\|\mu} = \frac{i}{2}\tau^3[\rho_{2\mu} - \frac{1}{2G_2}(gW_{3\mu} + g'B_\mu)], \tag{B.12}$$

$$V_{\omega 2\mu} - \frac{i}{G_{\omega 2}}\alpha_{\omega 2\|\mu} = \frac{i}{2} I_2[\omega_{2\mu} - \frac{1}{2G_{\omega 2}}(2Y_{Ll}g'B_\mu)]. \tag{B.13}$$

By substituting these expressions, we obtain :

$$\begin{aligned}
\mathcal{L}_S = \quad & \frac{1}{2}\{(\partial_\mu P^3)^2 + (\partial_\mu \Pi^3)^2 + (\partial_\mu \theta_8)^2\} + \frac{1}{2}\sum_{a=1}^{2}\{(\partial_\mu P^a)^2 + (\partial_\mu \hat{\Pi}^a)^2\} \\
+ \quad & \frac{1}{2}\sum_{a=1}^{2}(F_L \partial_\mu \hat{\Pi}^a + \sqrt{3}F_6 \partial_\mu P^a) gW_\mu^a + \frac{1}{2}(F_2\partial_\mu \Pi^3 + \sqrt{3}F_6\partial_\mu P^3)(gW_\mu^3 - g'B_\mu) \\
+ \quad & \frac{1}{8}(3F_6^2 + F_2^2)(gW_\mu^3 - g'B_\mu)^2 + \frac{1}{8}(3F_6^2 + F_L^2)(gW_\mu^a)^2 \\
- \quad & \frac{\sqrt{3}\beta_\perp}{4F_8F_2}\partial_\mu\theta_8\{\partial_\mu\Pi^3 + \frac{F_2}{2}(gW_\mu^3 - g'B_\mu)\} + \mathcal{L}'_{br},
\end{aligned} \tag{B.14}$$

$$\begin{aligned}
\mathcal{L}_V = \quad - \quad & \frac{1}{4}(\partial_{[\mu,}\rho_{6\nu]})^2 - \frac{1}{4}(\partial_{[\mu,}\omega_{6\nu]})^2 \\
- \quad & \frac{1}{4}(\partial_{[\mu,}\rho_{2\nu]})^2 - \frac{1}{4}(\partial_{[\mu,}\omega_{2\nu]})^2 - \frac{\alpha_V}{2}(\partial_{[\mu,}\rho_{2\nu]})(\partial_{[\mu,}\omega_{2\nu]}) \\
+ \quad & \frac{1}{2}M_{V6}^2\{\rho_{6\mu}^3 - \frac{\sqrt{3}}{2G_6}(gW_\mu^3 + g'B_\mu)\}^2 + \frac{1}{2}M_{V6}^2\{\rho_{6\mu}^a - \frac{\sqrt{3}}{2G_6}gW_\mu^a\}^2 \\
+ \quad & \frac{1}{2}M_{\omega 6}^2\{\omega_{6\mu} - \frac{\sqrt{3}}{2G_{\omega 6}}2g'Y_{Lq}B_\mu\}^2 \\
+ \quad & \frac{1}{2}M_{V2}^2\{\rho_{2\mu} - \frac{1}{2G_2}(gW_\mu^3 + g'B_\mu)\}^2 + \frac{1}{2}M_{\omega 2}^2\{\omega_{2\mu} - \frac{1}{2G_{\omega 2}}2g'Y_{Ll}B_\mu\}^2 \\
+ \quad & \frac{\beta_V}{2}\{\rho_{2\mu} - \frac{1}{2G_2}(gW_\mu^3 + g'B_\mu)\}\{\omega_{2\mu} - \frac{1}{2G_{\omega 2}}2g'Y_{Ll}B_\mu\},
\end{aligned} \tag{B.15}$$

$$\begin{aligned}
\mathcal{L}_A = \quad - \quad & \frac{1}{4}(\partial_{[\mu,}a_{6\nu]}^a)^2 - \frac{1}{4}(\partial_{[\mu,}a_{8\nu]})^2 \\
- \quad & \frac{1}{4}(\partial_{[\mu,}a_{2\nu]})^2 - \frac{\alpha_A}{2}(\partial_{[\mu,}a_{8\nu]})(\partial_{[\mu,}a_{2\nu]}) \\
- \quad & \frac{1}{4}\sum_{a=1}^{2}(\partial_{[\mu}a_{L\nu]}^a)^2
\end{aligned}$$

$$
\begin{aligned}
&+ \frac{1}{2}M_{A6}^2 \sum_{a=1}^{2}[a^a_{6\mu} + \frac{1}{\lambda_6 F_6}\{\partial_\mu P^a + \frac{\sqrt{3}F_6}{2} g W^a_\mu\}]^2 \\
&+ \frac{1}{2}M_{A6}^2 [a^3_{6\mu} + \frac{1}{\lambda_6 F_6}\{\partial_\mu P^3 + \frac{\sqrt{3}F_6}{2}(g W^3_\mu - g' B_\mu)\}]^2 \\
&+ \frac{1}{2}M_{A8}^2 [a_{8\mu} + \frac{1}{\lambda_8 F_8}\partial_\mu \theta_8]^2 \\
&+ \frac{1}{2}M_{A2}^2 [a^3_{2\mu} + \frac{1}{\lambda_2 F_2}\{\partial_\mu \Pi^3 + \frac{F_2}{2}(g W^3_\mu - g' B_\mu)\}]^2 \\
&+ \frac{\beta_A}{2}[a^3_{2\mu} + \frac{1+\delta}{\lambda_2 F_2}\{\partial_\mu \Pi^3 + \frac{F_2}{2}(g W^3_\mu - g' B_\mu)\}][a_{8\mu} + \frac{1+\delta'}{\lambda_8 F_8}\partial_\mu \theta_8] \\
&+ \frac{1}{2}M_{AL}^2 \sum_{a=1}^{2}[a^a_{L\mu} + \frac{1}{\lambda_L F_L}\{\partial_\mu \hat{\Pi}^a + \frac{F_L}{2} g W^a_\mu\}]^2 \\
&+ \frac{1}{8}\frac{\beta_A^2 M_{A2}^2 \delta^2}{M_{A2}^2 M_{A8}^2 - \frac{1}{4}\beta_A^2}\frac{1}{\lambda_2^2 F_2^2}\{\partial_\mu \Pi^3 + \frac{F_2}{2}(g W^3_\mu - g' B_\mu)\}^2 \\
&+ \frac{1}{8}\frac{\beta_A^2 M_{A8}^2 \delta'^2}{M_{A2}^2 M_{A8}^2 - \frac{1}{4}\beta_A^2}\frac{1}{\lambda_8^2 F_8^2}\partial_\mu \theta_8^2 \\
&- \frac{1}{2}\frac{\delta\delta'\beta_A M_{A2}^2 M_{A8}^2}{M_{A2}^2 M_{A8}^2 - \frac{1}{4}\beta_A^2}\frac{1}{F_2 F_8 \lambda_2 \lambda_8}\partial_\mu \theta_8\{\partial_\mu \Pi^3 + \frac{F_2}{2}(g W^3_\mu - g' B_\mu)\}, \qquad (\text{ B.16 })
\end{aligned}
$$

We fix the gauge into unitary gauge. This corresponds to the follwing replacement.

$$
\begin{aligned}
(g W^3_\mu - g' B_\mu) &\rightarrow (g W^3_\mu - g' B_\mu) - \frac{2}{3F_6^2 + F_2^2}(\sqrt{3}F_6 \partial_\mu P^3 + F_2 \partial_\mu \Pi^3), \qquad (\text{ B.17 }) \\
g W^a_\mu &\rightarrow g W^a_\mu - \frac{2}{3F_6^2 + F_L^2}(\sqrt{3}F_6 \partial_\mu P^a + F_2 \partial_\mu \hat{\Pi}^a), \qquad (\text{ B.18 })
\end{aligned}
$$

We also redefine the axial vector and left-handed vector mesons.

$$
\begin{aligned}
a^a_{6\mu} &\rightarrow a^a_{6\mu} - \frac{1}{\lambda_6 F_6}\partial_\mu P^a, \qquad (\text{ B.19 }) \\
a^3_{6\mu} &\rightarrow a^3_{6\mu} - \frac{1}{\lambda_6 F_6}\partial_\mu P^3, \qquad (\text{ B.20 }) \\
a_{8\mu} &\rightarrow a_{8\mu} - \frac{1}{\lambda_8 F_8}\partial_\mu \theta_8, \qquad (\text{ B.21 }) \\
a^3_{2\mu} &\rightarrow a^3_{2\mu} - \frac{1}{\lambda_2 F_2}\partial_\mu \Pi^3, \qquad (\text{ B.22 }) \\
a^a_{L\mu} &\rightarrow a^a_{L\mu} - \frac{1}{\lambda_L F_L}\partial_\mu \hat{\Pi}^a. \qquad (\text{ B.23 })
\end{aligned}
$$

Then, $\mathcal{L}_S$ and $\mathcal{L}_A$ are written in terms of physical degrees of freedom.

$$
\begin{aligned}
\mathcal{L}_S \;=\;& \frac{1}{2}\{(\partial_\mu\tilde{\Pi}^3)^2+(\partial_\mu\tilde{\Pi}^a)^2+(\partial_\mu\theta_8)^2\} \\
& -\frac{\sqrt{3}\beta_\perp}{4F_8F_2}\partial\theta_8\{\cos\chi_3\partial_\mu\tilde{\Pi}^3+\frac{F_2}{2}(gW^3_\mu-g'B_\mu)\} \\
& +\frac{1}{8}(3F_6^2+F_2^2)(gW^3_\mu-g'B_\mu)^2+\frac{1}{8}(3F_6^2+F_L^2)(gW^a_\mu)^2 \\
& +\mathcal{L}'_{br},
\end{aligned}
\qquad (\text{ B.24 })
$$

$$
\begin{aligned}
\mathcal{L}_A = \;-\;& \frac{1}{4}(\partial_{[\mu,}a^a_{6\nu]})^2-\frac{1}{4}(\partial_{[\mu,}a_{8\nu]})^2 \\
-\;& \frac{1}{4}(\partial_{[\mu,}a_{2\nu]})^2-\frac{\alpha_A}{2}(\partial_{[\mu,}a_{8\nu]})(\partial_{[\mu,}a_{2\nu]}) \\
-\;& \frac{1}{4}\sum_{a=1}^{2}(\partial_{[\mu,}a^a_{L\nu]})^2 \\
+\;& \frac{1}{2}M_{A6}^2\sum_{a=1}^{2}[a^a_{6\mu}+\frac{\sqrt{3}}{2\lambda_6}gW^a_\mu]^2 \\
+\;& \frac{1}{2}M_{A6}^2[a^3_{6\mu}+\frac{\sqrt{3}}{2\lambda_6}(gW^3_\mu-g'B_\mu)]^2 \\
+\;& \frac{1}{2}M_{A8}^2[a_{8\mu}]^2 \\
+\;& \frac{1}{2}M_{A2}^2[a_{2\mu}+\frac{1}{2\lambda_2}(gW^3_\mu-g'B_\mu)]^2 \\
+\;& \frac{\beta_A}{2}[a_{2\mu}+\frac{1}{2\lambda_2}(gW^3_\mu-g'B_\mu)][a_{8\mu}] \\
+\;& \frac{1}{2}M_{AL}^2\sum_{a=1}^{2}[a^a_{L\mu}+\frac{1}{2\lambda_L}gW^a_\mu]^2 \\
+\;& \frac{\beta_A\delta'}{2\lambda_8F_8}a_2^3\partial_\mu\theta_8+\frac{\beta_A\delta}{2\lambda_2F_2}a_8\{cos\chi_3\partial_\mu\tilde{\Pi}^3+\frac{F_2}{2}(gW^3_\mu-g'B_\mu)\} \\
+\;& \frac{1}{8}\frac{\beta_A^2M_{A2}^2\delta^2}{M_{A2}^2M_{A8}^2-\frac{1}{4}\beta_A^2}\frac{1}{\lambda_2^2F_2^2}\{cos\chi_3\partial_\mu\tilde{\Pi}^3+\frac{F_2}{2}(gW^3_\mu-g'B_\mu)\}^2 \\
+\;& \frac{1}{8}\frac{\beta_A^2M_{A8}^2\delta'^2}{M_{A2}^2M_{A8}^2-\frac{1}{4}\beta_A^2}\frac{1}{\lambda_8^2F_8^2}(\partial_\mu\theta_8)^2 \qquad (\text{ B.25 }) \\
-\;& \frac{1}{8}\frac{\delta\delta'\beta_A^3}{M_{A2}^2M_{A8}^2-\frac{1}{4}\beta_A^2}\frac{1}{F_2F_8\lambda_2\lambda_8}\partial_\mu\theta_8\{cos\chi_3\partial_\mu\tilde{\Pi}^3+\frac{F_2}{2}(gW^3_\mu-g'B_\mu)\},
\end{aligned}
$$

where,

$$\begin{pmatrix} \tilde{\Pi}^a \\ \tilde{P}^a \end{pmatrix} = \begin{pmatrix} cos\chi_L & -sin\chi_L \\ sin\chi_L & cos\chi_L \end{pmatrix} \begin{pmatrix} \hat{\Pi}^a \\ P^a \end{pmatrix}, \qquad (\text{B.26})$$

$$\begin{pmatrix} \tilde{\Pi}^3 \\ \tilde{P}^3 \end{pmatrix} = \begin{pmatrix} cos\chi_3 & -sin\chi_3 \\ sin\chi_3 & cos\chi_3 \end{pmatrix} \begin{pmatrix} \Pi^3 \\ P^3 \end{pmatrix}, \qquad (\text{B.27})$$

$$cos\chi_L = \frac{\sqrt{3}F_6}{\sqrt{3F_6^2 + F_L^2}}, \quad sin\chi_L = \frac{F_L}{\sqrt{3F_6^2 + F_L^2}}, \qquad (\text{B.28})$$

$$cos\chi_3 = \frac{\sqrt{3}F_6}{\sqrt{3F_6^2 + F_2^2}}, \quad sin\chi_3 = \frac{F_2}{\sqrt{3F_6^2 + F_2^2}}. \qquad (\text{B.29})$$

C Pseudo Nambu Goldstone boson sector

$$\begin{aligned} \mathcal{L}_S &= \frac{1}{2}\{(\partial_\mu\tilde{\Pi}^3)^2 + (\partial_\mu\tilde{\Pi}^a)^2 + (\partial_\mu\theta_8)^2\} \\ &+ \frac{1}{8}(3F_6^2 + F_2^2)(gW_\mu^3 - g'B_\mu)^2 + \frac{1}{8}(3F_6^2 + F_L^2)(gW_\mu^a)^2 \\ &-\frac{\sqrt{3}\beta_\perp}{4F_8F_2}\partial\theta_8\{\cos\chi_3\partial_\mu\tilde{\Pi}^3 + \frac{1}{2}(gW_\mu^3 - g'B_\mu)\} + \mathcal{L}'_{br}. \qquad (\text{C.1}) \end{aligned}$$

Now we are ready for giving the explicit form for $\mathcal{L}'_{br}$. Because $\tilde{\Pi}^3$, $\tilde{\Pi}^a$ and θ_8 are NGBs associated with broken global symmetry, we can introduce the follwing mass terms as $\mathcal{L}'_{br}$.

$$\mathcal{L}'_{br} = -\frac{1}{2}\begin{pmatrix} \tilde{\Pi}_3 & \theta_8 \end{pmatrix} \begin{pmatrix} m_3^2 & \frac{\beta_S}{2} \\ \frac{\beta_S}{2} & m_8^2 \end{pmatrix} \begin{pmatrix} \tilde{\Pi}_3 \\ \theta_8 \end{pmatrix} - \frac{1}{2}m_a^2\tilde{\Pi}_a^2 \qquad (\text{C.2})$$

The mass terms break the global symmetry without loss of $SU(2)\otimes U(1)$ gauge invariance and these NGBs become PNGBs.

D $O(p^2)$ terms in axial vector and left-handed vector sector

In this appendix, we show how to determine the $O(p^2)$ terms which consist of NGBs and PNGBs in this sector. As explained in the text, we add the $O(p^2)$ terms so that the incorporation of axial and left-handed vector mesons does not change the decay constants of NGBs and PNGBs. By doing so, ρ (T) parameter depends only on the parameters in $\mathcal{L}_s$. Though it is just the matter of the definition of the parameters of $O(p^2)$ terms in $\mathcal{L}_s$, our choise is convenient because the the parameters in $\mathcal{L}_s$ are directly related to physical quantities such as decay constants. Further ρ (T) parameter is independent of the parameters in $\mathcal{L}_A$ with the procedure adopted here. Let us discuss $0(p^2)$ terms which consist of $\alpha_{6\perp}$ and $\alpha_{L\perp}$ first. These terms are related to A_6 and A_L. The equations of motion of A_6 and A_L up to $O(p^2)$ are,

$$\begin{aligned} A_6 &= \frac{i}{\lambda_6}\alpha_{6\perp}, \\ A_L &= \frac{i}{\lambda_L}\alpha_{L\perp}. \end{aligned} \tag{D.1}$$

By substituting these into (44), we do not have $O(p^2)$ terms of NGBs and PNGBs. Therefore the $O(p^2)$ terms which are already present in (44) is enough. $O(p^2)$ terms which consist of $\alpha_{2\perp}$ and $\alpha_{8\perp}$ have more complicated coefficients as shown in (44). We have determined them in the follwing way. Let us focus on a part of $\mathcal{L}_A$

$$\begin{aligned} \mathcal{L}_0 = \quad & \frac{1}{2}M_{A_8}^2(a_{8\mu} - \frac{1}{\lambda_8}\hat{\alpha}_{8\perp\mu})^2 + \frac{1}{2}M_{A_2}^2(a_{2\mu} - \frac{1}{\lambda_2}\hat{\alpha}_{2\perp\mu})^2 \\ + \quad & \beta_A\frac{1}{2}(a_{2\mu} - \frac{(1+\delta)}{\lambda_2}\hat{\alpha}_{2\perp\mu})(a_{8\mu} - \frac{(1+\delta')}{\lambda_8}\hat{\alpha}_{8\perp\mu}) \\ \\ = \quad & \frac{1}{2}\left((a_2 - \tfrac{1}{\lambda_2}\hat{\alpha}_{2\perp}) \quad (a_8 - \tfrac{1}{\lambda_8}\hat{\alpha}_{8\perp}) \right) \begin{pmatrix} M_{A2}^2 & \frac{\beta_A}{2} \\ \frac{\beta_A}{2} & M_{A8}^2 \end{pmatrix} \begin{pmatrix} (a_2 - \frac{1}{\lambda_2}\hat{\alpha}_{2\perp}) \\ (a_8 - \frac{1}{\lambda_8}\hat{\alpha}_{8\perp}) \end{pmatrix} \\ & + \frac{1}{2}\left((a_2 - \tfrac{1}{\lambda_2}\hat{\alpha}_{2\perp}) \quad (a_8 - \tfrac{1}{\lambda_8}\hat{\alpha}_{8\perp}) \right) \begin{pmatrix} 0 & \beta_A\delta' \\ \beta_A\delta & 0 \end{pmatrix} \begin{pmatrix} -\frac{1}{\lambda_2}\hat{\alpha}_{2\perp} \\ -\frac{1}{\lambda_8}\hat{\alpha}_{8\perp} \end{pmatrix} \\ & + \frac{\beta_A}{2}\frac{\delta\delta'}{\lambda_2\lambda_8}\hat{\alpha}_{2\perp}\hat{\alpha}_{8\perp}, \end{aligned} \tag{D.2}$$

where we have used the follwing notation;

$$A_{2\mu} \;=\; i\frac{1}{2}\begin{pmatrix} 0 & \\ & \tau^3 \end{pmatrix} a_{2\mu}, \tag{D.3}$$

$$A_{8\mu} = i\frac{1}{4\sqrt{3}}\begin{pmatrix} I_6 & \\ & -3I_2 \end{pmatrix} a_{8\mu}, \tag{ D.4 }$$

$$\alpha_{2\perp\mu} = \frac{1}{2}\begin{pmatrix} 0 & \\ & \tau^3 \end{pmatrix} \hat{\alpha}_{2\perp\mu} \tag{ D.5 }$$

$$\alpha_{8\perp\mu} = \frac{1}{4\sqrt{3}}\begin{pmatrix} I_6 & \\ & -3I_2 \end{pmatrix} \hat{\alpha}_{8\perp\mu} \tag{ D.6 }$$

The equations of motion for a_2 and a_8 up to $O(p^2)$ are:

$$\begin{pmatrix} a_2 - \frac{1}{\lambda_2}\hat{\alpha}_{2\perp} \\ a_8 - \frac{1}{\lambda_8}\hat{\alpha}_{8\perp} \end{pmatrix} = \frac{1}{2}\frac{1}{M_{A2}^2 M_{A8}^2 - \frac{\beta_A^2}{4}} \begin{pmatrix} M_{A8}^2 & -\frac{\beta_A}{2} \\ -\frac{\beta_A}{2} & M_{A2}^2 \end{pmatrix} \begin{pmatrix} 0 & \beta_A \delta' \\ \beta_A \delta & 0 \end{pmatrix} \begin{pmatrix} \frac{1}{\lambda_2}\hat{\alpha}_{2\perp} \\ \frac{1}{\lambda_8}\hat{\alpha}_{8\perp} \end{pmatrix} \tag{ D.7 }$$

By substituting this into $\mathcal{L}_0$, we obtain;

$$\begin{aligned} \mathcal{L}_c = & \quad -\frac{1}{4}\frac{\beta_A^2 M_{A2}^2 \delta^2}{M_{A2}^2 M_{A8}^2 - \frac{1}{4}\beta_A^2}\frac{1}{\lambda_2^2} tr(\alpha_{2\perp\mu})^2 \\ & + \quad -\frac{1}{4}\frac{\beta_A^2 M_{A8}^2 \delta'^2}{M_{A2}^2 M_{A8}^2 - \frac{1}{4}\beta_A^2}\frac{1}{\lambda_8^2} tr(\alpha_{8\perp\mu})^2 \\ & + \quad -\frac{2}{\sqrt{3}}\frac{\delta\delta'\beta_A M_{A2}^2 M_{A8}^2}{M_{A2}^2 M_{A8}^2 - \frac{1}{4}\beta_A^2}\frac{1}{\lambda_2\lambda_8} tr(\alpha_{2\perp\mu}\tau^3\alpha_{8\perp\mu}) \end{aligned} \tag{ D.8 }$$

By subtracting $\mathcal{L}_c$ from $\mathcal{L}_0$, we obtain $O(p^2)$terms which consist of $\alpha_{2\perp}$ and $\alpha_{8\perp}$ in (44). These are the desired counter terms which kill the effect of axial vector mesons and left-handed vector mesons on ρ (T) parameter.

E S in QCD scale up technicolor model

For the completeness, we compute S_{theory} of QCD scale up technicolor model which is quoted in the introduction and (78). The QCD scale up technicolor model has $SU(2)_L \otimes SU(2)_R$ global symmetry and $N_{TC} = N_c = 3$. Therefore we only need to study the SU(2) subsector of one-family model. S in this model is given by;

$$S = 4\pi[\frac{1}{G_V^2} - \frac{1}{\lambda_A^2}]. \tag{ E.1 }$$

The G_V and λ_A are defined in the same way as G_6 and λ_6. G_V and λ_A are determined by $\rho \to \pi\pi$ and $a_1 \to \gamma\pi$ decays .

$$\begin{aligned} \Gamma_{\rho\pi\pi} &= \frac{1}{48\pi} m_\rho \left(\frac{m_\rho^2}{2G_V f_{QCD}^2}\right)^2 \left(1 - \frac{4m_\pi^2}{m_\rho^2}\right)^{\frac{3}{2}}, & \text{(E.2)} \\ \Gamma_{a\gamma\pi} &= \frac{\alpha}{24 f_{QCD}^2 G_A^2} \frac{(m_a^2 - m_\pi^2)^3}{m_a^3}, & \text{(E.3)} \end{aligned}$$

where f_{QCD} is the pion decay constant. By using the following values,

$$f_{QCD} = 93MeV, \quad m_\pi = 140MeV, \quad m_\rho = 768MeV, \quad m_a = 1260MeV,$$

$$\Gamma_{\rho\pi\pi} = 152MeV, \quad \Gamma_{a\gamma\pi} = 0.64MeV,$$

we obtain;

$$G_V^2 = 31.5, \quad G_A^2 = 106.$$

This leads to the following estimation of S_{theory} for the QCD scale up technicolor model which is quoted in the text.

$$S_{theory} = 0.28 \tag{ E.4 }$$

Because they are dimensionless quantities, the scaling relations between the parameters in QCD and that of the technicolor are given by,

$$G_6 = G_V, \quad \lambda_6 = \lambda_A. \tag{ E.5 }$$

This equalities hold if the underlying dynamics of the techniquark sector is the same as that of QCD.

References

[1] B. Holdom and J. Terning, *Phys. Lett.* **B247** (1990) 88 ;

M. Golden and L. Randall, *Nucl. Phys.* **B361** (1991) 3 ;

M. E. Peskin and T. Takeuchi, *Phys. Rev. Lett.* **65** (1990) 964 ; *Phys. Rev.* **D46** (1992) 381.

[2] R. Tanaka, Proceedings of XXVI International conference on High Energy Physics, (1992) 681, Dallas, Texas, Edited by J. R. Sanford.

[3] T. Appelquist and J. Terning, *Phys. Lett.* **B315** (1993) 139.

[4] S. Weinberg, *Phys. Rev.* **166**, (1968) 1568.

[5] M. Bando, T. Kugo, and K. Yamawaki,*Phys. Rep.***164** (1988) 217 ; and references therein.

[6] G. Ecker, A. Pich, and E. de Rafael, *Nucl. Phys.* **B321** (1989) 311, J. F. Donoghue, and B. R. Holstein, *Phys. Rev.* **D40** (1989) 2378.

[7] T.Onogi and Y.Yasui, *Mod. Phys. Lett.***A7**(1992) 785 ;

C. T. Hill, D. C. Kennedy, T. Onogi and H. L. Yu, *Phys. Rev.***D47** (1993) 2940.

Table Caption

- **Table(1)**: J,P,C,I and electric charge of the technimesons incorporated in the effective Lagrangian. In the third column, the numbers of corresponding technimesons are shown. In the fourth column, techniquark contents with the same J,P,C,I and electric charge are shown.

Figure Captions

- **Figure(1)**: The Feynman diagrams for the contribution to S in the techniquark sector.

- **Figure(2)**: The Feynman daigrams for the contribution to S in the technilepton sector.

- **Figure(3)**: The Feynman diagram for the contribution to $\delta\Pi_{11}$ part of U due to the left-handed vector meson.

- **Figure(4)**: $S < 0$ ($S > 0$) region with vector(1^{--}) dominance assumption for $\alpha_V = 1$. G_6, G_2 and $G_{\omega 2}$ are coupling constants associated with vector mesons. (See (31) in the text.)

	J^{pc}	$Number$		$Isospin$	$charge$
P^α	0^-	35	$\bar{Q}\gamma_5 T^\alpha Q$		
Π^3	0^{-+}	1	$\bar{L}\gamma_5\tau^3 L$	1	0
θ_8	0^{-+}	1	$\bar{Q}\gamma_5 Q - 3\bar{L}\gamma_5 L$	0	0
$\Pi^\pm$	0	2	$\bar{L}\frac{1}{2}(1-\gamma_5)\tau^\pm L$	1	±1
$V^\alpha_{6\mu}$	1^-	35	$\bar{Q}\gamma_\mu\lambda^\alpha Q$		
$\omega_{6\mu}$	1^{--}	1	$\bar{Q}\gamma_\mu Q$	0	0
$\rho_{2\mu}$	1^{--}	1	$\bar{L}\gamma_\mu\tau^3 L$	1	0
$\omega_{2\mu}$	1^{--}	1	$\bar{L}\gamma_\mu L$	0	0
$A^\alpha_{6\mu}$	1^+	35	$\bar{Q}\gamma_\mu\gamma_5 T^\alpha Q$		
$A_{8\mu}$	1^{++}	1	$\bar{Q}\gamma_\mu\gamma_5 Q - 3\bar{L}\gamma_\mu\gamma_5 L$	0	0
$A_{2\mu}$	1^{++}	1	$\bar{L}\gamma_\mu\gamma_5\tau^3 L$	1	0
$A^\pm_{L\mu}$	1	2	$\bar{L}\gamma_\mu\frac{1}{2}(1-\gamma_5)\tau^\pm L$	1	±1

Table 1



Fig.1



Fig.2



Fig.3



Fig.4

This figure "fig1-1.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9403343v2

This figure "fig2-1.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9403343v2

This figure "fig1-2.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9403343v2

This figure "fig2-2.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9403343v2

This figure "fig1-3.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9403343v2

This figure "fig1-4.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9403343v2

This figure "fig1-5.png" is available in "png" format from:

http://arXiv.org/ps/hep-ph/9403343v2